

FORM PTO-1390
(REV 10-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

GJE-230A

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

09/700821INTERNATIONAL APPLICATION NO.
PCT/GB00/03851INTERNATIONAL FILING DATE
October 6, 2000PRIORITY DATE CLAIMED
October 8, 1999

TITLE OF INVENTION
Supported Homogeneous Catalysts For Selective Hydrogenation

APPLICANT(S) FOR DO/EO/US

Mark Joseph Burk, Arne Gerlach

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)).
4. ☐ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Copy of Form PCT/RO/101
 Copy of Form PCT/RO/144
 Copy of Form PCT/RO139

COPY

U.S. APPLICATION NO. (if known, see 37 CFR 1.51) 09/700821		INTERNATIONAL APPLICATION NO. PCT/GB00/03851		ATTORNEY'S DOCKET NUMBER GJE-230A	
--	--	---	--	--------------------------------------	--

17. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1000.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00

International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00

International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total claims	14 - 20 =	0	X \$18.00	\$
Independent claims	1 - 3 =	0	X \$80.00	\$
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00	\$
TOTAL OF ABOVE CALCULATIONS =				\$ 860.00

☐ Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.

SUBTOTAL =

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(f)).

TOTAL NATIONAL FEE =

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property

TOTAL FEES ENCLOSED =

	Amount to be refunded:	\$
	charged:	\$

COPY

a. ☐ A check in the amount of \$_____ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 19-0065 in the amount of \$ 860.00 to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 19-0065. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Mr. David R. Saliwanchik
Saliwanchik, Lloyd & Saliwanchik
A Professional Association
2421 N.W. 41st Street, Suite A-1
Gainesville, FL 32606-6669

COPY

David Saliwanchik

SIGNATURE

David R. Saliwanchik

NAME

31,794

REGISTRATION NUMBER

09/700821

532 Rec'd PCT/PTO 20 NOV 2000

November 20, 2000

Patent Application
Docket No. GJE-230A

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) : Mark Joseph Burk and Arne Gerlach
Docket No. : GJE-230A
For : Supported Homogeneous Catalysts For Selective Hydrogenation

Box PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Please amend the above-identified application as follows:

In the Claims

Please amend the following claims:

Claim 2, line 1: Please delete "A" and insert --The--

Claim 3, line 1: Please delete "A" and insert --The--

Claim 4, line 1: Please delete "A" and insert --The--

Claim 5, line 1: Please delete "A" and insert --The--

Claim 6, line 1: Please delete "A" and insert --The--

Claim 6, line 1: Please delete "any preceding claim" and insert --claim 1--

Claim 7, line 1: Please delete “A” and insert --The--

Claim 8, line 1: Please delete “A” and insert --The--

Claim 8, line 1: Please delete “any preceding claim” and insert --claim 1--

Claim 9, line 1: Please delete “A” and insert --The--


Claim 10, line 1: Please delete “A” and insert --The--.

Claim 10, line 1: Please delete “any preceding claim” and insert --claim 1--

Claim 11, line 1: Please delete “A” and insert --The--

Claim 12 (amended):

[Use of a catalyst according to any preceding claim, in a] A process of hydrogenating an aldehyde to produce the corresponding primary alcohol wherein said process utilizes a supported catalyst comprising a cationic rhodium (I) complex of the formula



wherein R¹ and R² are the same or different hydrocarbon groups of up to 30 C atoms, or R¹ and R² are linked to form a ring, and a heterogenous support medium that provides anionic binding sites.

Claim 13, line 1: Please delete "Use" and insert --The process--

Claim 14, line 1: Please delete "Use" and insert --The process--

Claim 14, line 1: Please delete "or claim 13"

The Commissioner is hereby authorized to charge any fees under 37 CFR 1.16 or 1.17 as required by this paper to Deposit Account 19-0065.

Respectfully submitted,



David R. Saliwanchik

Patent Attorney

Registration No. 31,794

Phone No.: 352-375-8100

Address : 2421 N.W. 41st Street

Suite A-1

Gainesville, FL 32606

DRS/la

SUPPORTED HOMOGENEOUS CATALYSTS FOR
SELECTIVE HYDROGENATION

Field of the Invention

5 This invention relates to a supported catalyst and to its use, e.g. in the efficient and selective hydrogenation of aldehydes to alcohols.

Background of the Invention

Both homogeneous and heterogeneous catalysts are known, as are their respective advantages and disadvantages. One way of combining the features of both is to immobilise
10 or tether a homogeneous catalyst to a polymeric or inorganic solid support. An undesirable aspect of this strategy is that the heterogenised ligand systems often are very tedious and/or expensive to prepare. Another problem is that polymer-supported homogeneous catalysts frequently have reduced catalytic activities and selectivities relative to the unsupported homogeneous analogues. Upon attempted reuse, the activities and
15 selectivities of these catalysts are often reduced further. Finally, many immobilised homogeneous catalysts suffer from a high degree of metal loss from the support (leaching) during use; see, for example, Lindner, *et al*, *Angew. Chemie Int. Ed.* **1999**, 38, 2155.


Aldehyde reduction often is a desirable step in obtaining valuable alcohol products from inexpensive starting materials (e.g., alkenes, hydrogen and carbon monoxide in the
20 case of hydroformylation). Despite the importance of aldehyde reduction in organic chemistry, surprisingly few generally applicable manufacturing methods are available for this transformation. Hydride reducing agents (e.g. LiAlH_4 or NaBH_4) are widely used, but are moisture-sensitive reagents that are not economically attractive for commercial procedures since they are employed in stoichiometric quantities. Moreover, their use
25 requires tedious work-up procedures and generates substantial quantities of waste (boron or aluminium salts).

Numerous heterogeneous catalysts, such as PtO_2 , Raney Ni, and Pd/C, can catalyse the reduction of specific aldehydes. However, heterogeneous catalysts are often intolerant of various organic groups such as divalent sulfides. Moreover, other sensitive groups such
30 as nitro, oxime, ketone, aryl halide or benzyloxy, also are reduced. Another problem encountered when reducing aromatic aldehydes using heterogeneous catalysts is that the

product may be further reduced to a methyl substituent. For example, heterogeneous hydrogenation of benzaldehyde often affords toluene.

Very few practical homogeneous systems efficiently catalyse aldehyde hydrogenation. Problems often encountered include low reaction rates and/or catalyst deactivation due to aldehyde decarbonylation processes.

The use of cationic rhodium catalysts for aldehyde hydrogenation has been reported by Tani *et al*, *Chem. Lett.* **1982**, 261, and by Burk *et al*, *Tetrahedron Lett.* **1994**, 35, 4963. Results suggest that the achievement of high efficiency in rhodium-catalysed aldehyde hydrogenation requires the use of electron-rich (dialkyl- or trialkyl-substituted) chelating phosphine ligands, but these tend to be very air-sensitive and are not suitable for industrial manufacture. Burk *et al*. describes an electron-rich, yet air-stable crystalline ligand, 1,1'-bis(diisopropylphosphino)ferrocene (DiPFc) **1** ($R^1 = R^2 = i\text{-Pr}$). The homogeneous rhodium catalyst (DiPFc-Rh) also is stable to oxygen and has been shown to hydrogenate a limited set of aldehydes with high reaction rates.



A new method of anchoring certain rhodium catalysts to solid supports has recently been described by Augustine *et al*, *Chem. Comm.* **1999**, 1257. This simple procedure involves treating a readily available solid material (silica, alumina, carbon, etc.) with a heteropolyacid such as phosphotungstic acid, followed by addition of an appropriate catalyst precursor complex. Immobilised catalysts formed in this fashion were reported to serve as active and reusable catalysts for alkene hydrogenation.

Summary of the Invention

According to the present invention, an immobilised homogeneous catalyst is useful, *inter alia*, for the efficient and chemoselective hydrogenation of aldehydes. The catalyst system is based upon homogeneous rhodium complexes bearing phosphines of formula **1**, wherein R^1 and R^2 are independently the same or different hydrocarbon substituents, e.g.

alkyl, substituted alkyl, arylalkyl or aryl, of up to 30 C atoms, or R^1 and R^2 are linked to form a ring. Preferably, R^1 and R^2 are alkyl groups and more preferably identical alkyl groups. The solid support provides anionic binding sites.

The utility of the novel catalyst is surprising, for various reasons. Firstly, it was not evident that the complex could be supported. Secondly, its activity for aldehyde hydrogenation is good especially given the acidic nature of the support. Further, the immobilised hydrogenation catalyst can be effectively recovered and re-used.

Description of the Invention

Solid supports that are effective for use in the invention are those providing anionic binding sites. The support may or may not be modified with a heteropolyacid anchoring agent. The support medium is preferably an oxide such as alumina, silica, carbon, montmorillonite, etc., and is preferably modified with a heteropolyacid. The heteropolyacid is preferably of the Keggin type, e.g. phosphotungstic acid, phosphomolybdic acid or silicotungstic acid. Alternatively, an anionic exchange resin such as poly *para*-toluenesulfonic acid or Nafion in its acidic or anionic form may be used. For example, the support medium is a cation exchange resin containing sulphonic acid groups $-\text{SO}_3^- \text{X}^+$, wherein X^+ is a proton or any other exchangeable cation. A preferred cation exchange resin is a tetrafluoroethylene-perfluoro(vinyl ether sulfonate) copolymer.

Many different types of aldehydes, e.g. of formula RCHO , wherein R is an organic group up to 30 C atoms, may be hydrogenated to give RCH_2OH , using the novel catalyst. The aldehyde substrate may possess a range of different functional groups that either inhibit or react with commonly employed heterogeneous catalysts. Due to the acidic nature of the supports used in the immobilisation of the homogeneous catalyst, a non-standard solvent mixture may be required. The use of an alcohol/water mixture, and particularly an isopropanol/water mixture, is preferred, so that the hydrogenation reaction proceeds to completion. In particular, acetal formation can be minimised or avoided. The immobilised catalyst system may be recovered by simple filtration and re-used in subsequent reactions.

In addition to the hydrogenation of aldehydes, a catalyst of the invention may also be used for hydrogenation of other unsaturated groups. For example, unsaturated functionality such as the carbon-carbon double bond of alkenes, the carbon-carbon triple bond of alkynes, the carbon-oxygen double bond of ketones and the carbon-nitrogen


double bond of substrates such as N-acylhydrazones may be hydrogenated using these catalyst systems.

Some aldehydes are quite temperature-sensitive and decompose as the temperature is increased much above room temperature. In such cases, the ability to perform the hydrogenation at mild temperatures is vital. Increasing the temperature may increase the reaction rate, but the novel catalyst may be performed over a broad temperature range of -30°C to $+150^{\circ}\text{C}$. The preferred temperature is in the range 0°C to 60°C .

Performing reactions under low pressure is often preferred for manufacturing due to the fact that high-pressure equipment is more costly to purchase and operate. An important advantage of this invention is that the catalyst can perform effectively under both high and low hydrogen pressures, e.g. over the range of 1 to 100 atmospheres (100 - 10000 kPa). Increasing the pressure may increase the reaction rates. The preferred pressure range will depend on the process being operated and the desired reaction rates.

Heterogenised rhodium catalyst systems bearing 1,1'-bis(dialkylphosphino)ferrocene ligands **1** may be prepared via various procedures. By way of representative example, an immobilised catalyst may be formed by mixing neutral alumina with phosphotungstic acid in methanol, followed by the addition of the catalyst precursor $[(\text{COD})\text{Rh}(\text{DiPFC})]^+\text{BF}_4^-$ (see Scheme 1). After allowing the mixture to stir for a specified period, the rhodium complex is completely absorbed onto the solid support. The tethered catalyst is then filtered, washed with methanol, and employed directly in catalysis. The mechanism of absorption and the exact nature of the tethered complex are unclear.

Scheme 1.



The DiPFc-Rh catalyst prepared as in Scheme 1 has been tested for effectiveness in the hydrogenation of a range of different multifunctional aldehydes. These studies were aimed at demonstrating the combined properties of high catalytic efficiency under mild conditions, selectivity in the reduction process, and tolerance of the catalyst to certain functionality. The robust nature of the catalyst system also was important. Moreover, comparisons have been made with commonly employed heterogeneous catalysts such as palladium on carbon, platinum oxide, and palladium on barium sulfate.

The following Examples illustrate the invention.

10 Example 1

Preparation of [(DiPFc)Rh(COD)]X on modified silica

A solution of phosphotungstic acid (PTA, 288 mg, 0.1 mmol, 1.0 eq.) in 25 ml degassed methanol was added dropwise to a vigorously stirred (overhead stirrer was used to minimise grinding) suspension of 4.00 g silica (Silica gel 60 for flash chromatography (Fluka), particle size 0.035-0.070 mm (220-440 mesh ASTM, activity according to Brockmann and Schroder: 2-3) in 30 ml of degassed methanol under nitrogen. The resulting mixture was stirred for 1 hour at room temperature. Subsequently, a solution of [(DiPFc)Rh(COD)]BF₄ (64 mg, 0.09 mmol, 0.9 eq.) in 10 ml degassed methanol was dripped to the vigorously stirred slurry of the activated silica. Stirring was continued for 4.5 hours at room temperature. After solvent evaporation the remaining solid was placed in a Soxhlet apparatus and continuously extracted with degassed methanol under nitrogen for 16 hours. The orange silica powder was isolated, dried and stored under nitrogen as a precaution. Yield: 3.64 g (86%).

The alumina supported DiPFc-Rh catalyst was prepared by an analogous protocol to that outlined above.

Example 2

General Hydrogenation Procedure

All reactions were carried out in a 50 ml Parr micro-reactor modified with an injection septum and valve. The micro reactor was used in connection with a suitable glass liner. The solvent (2-propanol/water mixture; 1:1 v/v) was deoxygenated by bubbling nitrogen through it for 3 hours while stirring. The hydrogenation substrate and the immobilised catalyst were added to a 50 ml glass liner, which was then immediately placed

in a 50 ml Parr pressure vessel. This was then sealed and purged with hydrogen (5 pressurisation (690 kPa)/release cycles). Degassed solvent (2-propanol/water; 1:1 v/v) was then added *via* cannula, the reactor purged again with hydrogen (5 pressurisation (690 kPa)/release cycles), charged to the initial hydrogen pressure (690 kPa) and vigorously stirred at a constant temperature (ambient temperature or heating bath). After an allocated period of time (hydrogen uptake was monitored) hydrogen pressure was released, and the reaction mixture was filtered (separation from the supported catalyst). The filtrate was then extracted several times with dichloromethane. The combined organic extracts were dried over sodium sulfate, filtered and evaporated. The product distribution of the crude product mixture was determined by ^1H -NMR spectroscopy and was compared with authentic samples of all products. In cases where the formation of water-soluble or volatile products was likely, the hydrogenation mixture also was analysed *via* HPLC prior to extractive work-up.

Thus, experiments were performed under a standard set of mild reaction conditions: conversion to product = 100%, hydrogen pressure = 690 kPa, temperature = 20°C , reaction time = 16 h, mol aldehyde/mol Rh = 300-500 (based upon analysis of Rh content), concentration = 0.1 M, solvent: 2-propanol/water (1:1 v/v). Analytical procedures and results are given in Table 1.

The results show that the catalyst was robust and would operate effectively under very mild reaction conditions. This is demonstrated by the fact that all experiments listed in Table 1 were conducted using catalyst that was stored under an atmosphere of air for a period of ten months. This immobilised homogeneous catalyst allowed complete hydrogenation of each aldehyde listed to afford exclusively the desired alcohol product in high yield. The results further reveal that aldehydes bearing either alkyl substituents ($\text{R} = \text{alkyl}$) or aromatic substituents ($\text{R} = \text{aryl}$) may be reduced with equal facility. Functional groups that are reduced by most common heterogeneous catalysts, including aryl halide, nitro, and benzyloxy, were not reduced.

In contrast, common heterogeneous catalysts invariably yielded mixtures of products due to low chemoselectivity in reduction of the aldehyde carbonyl group (substrates 7-9, 13). In all cases using the heterogeneous catalysts, milder than normal reaction conditions were employed in an effort to achieve some level of selectivity in the reduction process. This strategy provided the best advantage to the heterogeneous

















catalyst, but resulted in incomplete conversion of starting aldehyde in some cases. In all such cases, driving the reactions further to allow complete conversion of substrate led to lower selectivities.














Importantly, the novel catalyst displayed broad tolerance to various organic functionalities, including sulphur-containing groups, with no apparent diminution of catalytic rates. Sulphur functionality is notoriously detrimental to most heterogeneous catalysts, leading to serious levels of catalyst inhibition. This point was amply demonstrated in experiments 10-13. Of particular note is the successful hydrogenation of substrates 10 and 12, which contain non-aromatic sulfide groups.

In addition to a catalyst with an alumina solid support, identical results were achieved using a DiPFc-Rh catalyst anchored to silica in the fashion described above in Scheme 1 (see hydrogenation results involving substrate 12). The use of a silica support offers significant practical advantages since this immobilised catalyst system is more readily handled and removed from the reaction mixtures.

One advantage of an immobilised catalyst is the potential to remove it completely from the reaction mixture through filtration, and also to reuse the catalyst in subsequent processes. This is demonstrated by performing 4 successive hydrogenations involving 2-thiophene carboxaldehyde (substrate 11). This particular aldehyde bears sulfur functionality, which should test the robustness of the immobilised catalyst in the presence of potential coordinating groups. In each case the hydrogenation was performed under conditions described in Table 1. After allowing the reaction to stir for 6 h (hydrogen uptake was monitored), a small sample was removed, and complete conversion to alcohol product was confirmed by ^1H NMR spectroscopy. The entire solution phase containing the product then was removed by syringe, the catalyst was washed twice with fresh solvent, and a subsequent aliquot of hydrogenation substrate in 2-propanol/water was added. The immobilised catalyst was used successfully for four catalytic cycles, and complete conversion to the corresponding alcohol product was observed after each run. No reduction of catalytic activity was noted over the four cycles.

Table 1. Selective Aldehyde Hydrogenations


Catalyst ^[a]		Product Distribution ^[b]			
 6	DiPFc-Rh on alumina	 CHO	 CH ₂ OH		
		n.d. ^[c]	100%		
 7	DiPFc-Rh on alumina	 CHO Br	 CH ₂ OH Br	 CHO	 CH ₂ OH
	dppb-Rh on silica	n.d.	100%	n.d.	n.d. ^[e]
	Platinum oxide	93%	7%	n.d.	n.d.
	Pd on carbon	1%	90%	<1%	8%
	Pd on carbon ^[d]	80%	n.d.	20%	n.d. ^[e]
	Pd on barium sulfate ^[f]	n.d.	n.d.	n.d.	100% ^[e]
 8	DiPFc-Rh on alumina	 CHO NO ₂	 CH ₂ OH NO ₂	nitro-reduced products	
	Pd on carbon	n.d.	100%	n.d.	
		58%	0%	42%	
 9	DiPFc-Rh on alumina	 CHO OBn	 CH ₂ OH OBn	 CHO OH	 CH ₂ OH OH
	Pd on carbon	n.d.	100%	n.d.	n.d.
		n.d.	n.d.	40%	60%

Catalyst ^[a]		Product Distribution ^[b]		
 10	DiPFc-Rh on alumina	n.d.	100%	
	Pd on carbon ^[i]	100%	n.d.	
	Platinum oxide ^[j]	>98%	<2%	
		 		
 11	DiPFc-Rh on alumina	n.d.	100%	
	Pd on carbon ^[i]	92%	8%	
		 		
 12	DiPFc-Rh on alumina	n.d.	100%	
	DiPFc-Rh on silica	n.d.	100%	
	DiPFc-Rh on silica ^[i]	n.d.	100%	
	DiPFc-Rh on silica ^[k]	69%	31%	
	dppb-Rh on silica	100%	0%	
	Platinum oxide ^[j]	3%	97%	
	Pd on carbon ^[i]	95%	5%	
		 		
 13	DiPFc-Rh on alumina	40%	60%	n.d.
	Pd on carbon	17%	n.d.	83%
		  		

[a] Hydrogenation conditions for DiPFc-Rh on alumina or silica: 100 psi hydrogen, S/C > 300, 0.1 molar in 2-propanol - water (1:1), room temperature, overnight; conditions for dppb-Rh on silica: 100 psi hydrogen, S/C > 200, 0.1 molar in 2-propanol - water (1:1), room temperature, overnight; conditions for platinum oxide: 1 bar hydrogen, 5 mg catalyst per mmol substrate, 0.1 molar 2-propanol - water (1:1), room temperature, 30 mins; conditions for palladium on carbon (10%): 1 bar hydrogen, 5 mg catalyst per mmol substrate, 0.1 molar in 2-propanol - water (1:1), room temperature, 30 mins. [b] Determined by ^1H -NMR analysis after extraction of the crude reaction mixture with dichloromethane, drying (sodium sulfate) and evaporation. [c] Not detected by ^1H -NMR, GC or HPLC. [d] Reaction conditions: 100 psi hydrogen, 10 mg palladium on carbon (10%) per mmol substrate, room temperature, 1 hour. [e] Determined by HPLC analysis of the crude reaction mixture. [f] Reaction conditions: 1 bar hydrogen, 5 mg palladium on barium sulfate (5%) per mmol substrate, room temperature, 30 mins. [g] Detected by HPLC analysis of the crude reaction mixture. [h] Purchased from Fluka as a technical mixture of various amounts of monomers and oligomers. [i] Reaction conditions: 100 psi hydrogen, 10 mg palladium on carbon (10%) per mmol substrate, room temperature, 16 hours. [j] Reaction conditions: 100 psi hydrogen, S/C > 500, 60°C, 20 hours. [k] Reaction conditions: 100 psi hydrogen, S/C > 1000, 60°C, 24 hours. [l] Reaction conditions: 100 psi hydrogen, 10 mg platinum oxide per mmol substrate, room temperature, 16 hours.

CLAIMS

1. A supported catalyst comprising a cationic rhodium(I) complex of the formula




- wherein R^1 and R^2 are the same or different hydrocarbon groups of up to 30 C atoms, or R^1 and R^2 are linked to form a ring, and a heterogeneous support medium that provides anionic binding sites.
2. A catalyst according to claim 1, wherein the support medium comprises a heteropolyacid anchoring agent.
3. A catalyst according to 2, wherein the heteropolyacid is of the Keggin type.
4. A catalyst according to claim 3, wherein the heteropolyacid is phosphotungstic acid, phosphomolybdic acid or silicotungstic acid.
5. A catalyst according to claim 4, wherein the heteropolyacid is phosphotungstic acid.
6. A catalyst according to any preceding claim, wherein the support medium comprises an oxide selected from alumina, silica, titania, lanthana, zeolites and clays.
7. A catalyst according to claim 6, wherein the metal oxide is alumina.
8. A catalyst according to any preceding claim, wherein the support medium is a cation exchange resin containing sulphonic acid groups $-SO_3^- X^+$, wherein X^+ is a proton or any other exchangeable cation.
9. A catalyst according to claim 8, wherein the cation exchange resin is a tetrafluoroethylene-perfluoro(vinyl ether sulfonate) copolymer.
10. A catalyst according to any preceding claim, wherein R^1 and R^2 are each an alkyl group.
11. A catalyst according to claim 10, wherein $R^1 = R^2 = i\text{-Pr}$.
12. Use of a catalyst according to any preceding claim, in a process of hydrogenating an aldehyde to produce the corresponding primary alcohol.

- 5

ABSTRACT**SUPPORTED HOMOGENEOUS CATALYSTS FOR
SELECTIVE HYDROGENATION**

5

A supported catalyst comprises a cationic rhodium(I) complex of the formula



10

wherein R^1 and R^2 are the same or different hydrocarbon groups of up to 30 C atoms, or
15 R^1 and R^2 are linked to form a ring, and a heterogeneous support medium that provides anionic binding sites.

Such a complex is particularly useful as a catalyst in a process of hydrogenating an aldehyde to produce the corresponding primary alcohol.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C 1001 and that such willful false statements may jeopardise the validity of the application or any patent issued thereon.

1-00
Full name of sole or
First Inventor

Mark Joseph BURK

Inventor's signature

Mark Joseph Burk

Residence address

California, USA

Post Office address

1610 Nantucket Circle, Apartment 115, Santa Clara, CA CA
95054, USA

Country of Citizenship United Kingdom

Date of signature 1 November 2000

2-00
Full name of
Second Inventor

Arne GERLACH

Inventor's signature

Arne Gerlach

Residence address

Cambridge, United Kingdom GBX

Post Office address

c/o Chirotech Technology Limited, Cambridge Science Park,
Milton Road, Cambridge CB4 0WG, United Kingdom

Country of Citizenship Germany

Date of signature 11 Nov. 2000

DECLARATION AND POWER OF ATTORNEY

As a below-named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of subject matter which is claimed and for which a patent is sought on an invention entitled
SUPPORTED HOMOGENEOUS CATALYSTS FOR SELECTIVE HYDROGENATION

the specification of which ☐ is attached hereto or

☐ was filed on 06 OCT 2000 as United States Application Number or PCT International Application Number PCT/GB00/03851 and was amended on (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56. I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for a patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed:

Prior Foreign Application Number(s)	Country	Foreign Filing Date	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
9923952.7	GB	08 OCT 1999	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:
 David R. Saliwanchik, Reg. 31,794; Jeff Lloyd, Reg. 35,589; Doran R. Pace, Reg. 38,261; Christine Q. McLeod, Reg. 36,213; Jay M. Sanders, Reg. 39,355; James S. Parker, Reg. 40,119 and Jean E. Kyle, Reg. 36,987; Frank C. Eisenschenk, Reg. 45,332; Seth M. Blum, Reg. 45,489

Direct all correspondence to:
Saliwanchik, Lloyd & Saliwanchik
2421 N.W. 41st Street, Suite A-1
Gainesville, FL 32606-6669
 USA

09/00/00 11:00:00

9